



US006011049A

# United States Patent [19]

## Whitcomb

[11] **Patent Number:** **6,011,049**  
[45] **Date of Patent:** **Jan. 4, 2000**

[54] **COMBINATIONS FOR DIABETES**

[75] Inventor: **Randall Wayne Whitcomb**, Ann Arbor, Mich.

[73] Assignee: **Warner-Lambert Company**, Morris Plains, N.J.

[21] Appl. No.: **09/189,132**

[22] Filed: **Nov. 9, 1998**

### Related U.S. Application Data

[63] Continuation-in-part of application No. 08/970,057, Nov. 13, 1997, Pat. No. 5,859,037

[60] Provisional application No. 60/038,224, Feb. 19, 1997.

[51] **Int. Cl.<sup>7</sup>** ..... **A61K 31/44**; A61K 31/425;  
A61K 31/175; A61K 31/155

[52] **U.S. Cl.** ..... **514/369**; 514/342; 514/593;  
514/635; 514/866

[58] **Field of Search** ..... 514/342, 369,  
514/593, 635, 866

[56] **References Cited**

#### FOREIGN PATENT DOCUMENTS

0749751 12/1996 European Pat. Off. .  
0753298 1/1997 European Pat. Off. .  
9609823 4/1996 WIPO .

#### OTHER PUBLICATIONS

PCT International Search Report, PCT/US97/21996.  
Oakes, et al., *Diabetes*, vol. 13:1203, 1994, A New Antidiabetic Agent, BRL 49653, Reduces Lipid Availability and Improves Insulin Action and Glucoregulation in the Rat.  
Hulin, et al, *Current Pharmaceutical Design* 2:85-102, 1996, The Glitazone Family of Antidiabetic Agents.

Groop, M.D., *Diabetes Care*, 15:737-754, 1992, Sulfonylureas in NIDDM.

Nakano, et al., *CS-045*, Clinical Evaluation of a New Oral Hypoglycemic Drug, CS-045, on Daily Profile of Blood Glucose in Patients with Non-Insulin Dependent Diabetes Mellitus (1993).

*Company News On-Call*, 1-3, 1997 Rezulin® (troglitazone) Receives FDA Marketing Clearance for use as Either Initial or Combination Therapy for Type 2 Diabetes.

Spencer, et al., *Drugs*, vol. 54:89-101, 1997, Troglitazone. Yamasaki, et al., *The Tokohu Journal of Experimental Medicine*, vol. 183:173-183, 1997, Pioglitazone (AD-4833) Ameliorates Insulin Resistance in Patients with NIDDM.

Iwamoto, et al., *Diabetic Medicine*, vol. 13:365-370, 1996, Effect of Combination Therapy of Troglitazone and Sulphonylureas in Patients with Type 2 Diabetes Who Were Poorly Controlled by Sulphonylurea Alone.

Bressler & Johnson., *Drugs & Aging*, vol. 9:418-437, 1996, Oral Antidiabetic Drug Use in the Elderly.

Okaka, et al., *Chemical Abstracts*, vol. 126, No. 20, 1997, Abstract No. 258925, Antidiabetic Effects of Pioglitazone HCL Alone or in Combination with Insulin or Sulphonylurea in Diabetic Animals.

De Souza, et al., *Diabetes*, vol. 44:984-991, 1995, Insulin Secretory Defect in Zucker fa/fa Rats Improved by Ameliorating Insulin Resistance.

*Primary Examiner*—Kimberly Jordan

*Attorney, Agent, or Firm*—Charles W. Ashbrook

[57] **ABSTRACT**

Combinations of a glitazone antidiabetic agent and a biguanide antidiabetic agent, and optionally a sulfonylurea antidiabetic agent, are useful for treating diabetes mellitus and improving glycemic control.

**16 Claims, 12 Drawing Sheets**